

QUOTIENT SPACES DETERMINED BY ALGEBRAS OF CONTINUOUS FUNCTIONS

ALDO J. LAZAR

ABSTRACT. We prove that if X is a locally compact σ -compact space then on its quotient, $\gamma(X)$ say, determined by the algebra of all real valued bounded continuous functions on X , the quotient topology and the completely regular topology defined by this algebra are equal. It follows from this that if X is second countable locally compact then $\gamma(X)$ is second countable locally compact Hausdorff if and only if it is first countable. The interest in these results originated in [1] and [7] where the primitive ideal space of a C^* -algebra was considered.

1. INTRODUCTION

The primitive ideal space, $\text{Prim}(A)$, of a C^* -algebra A with its hull-kernel topology has some pleasant properties: it is a locally compact Baire space, see [5, Corollary 3.3.8 and Corollary 3.4.13]. However, no better separation property than T_0 can be expected in general. The absence of the Hausdorff separation property in $\text{Prim}(A)$ justified a study in [11] of the collection of all the closed limit sets of a topological space. It should be added that the closed limit subsets of $\text{Prim}(A)$ correspond, by one of the bijections detailed in [5, Proposition 3.2.1], to some distinguished ideals of A . An investigation of the topologies on the class of these ideals that mirror topologies on the collection of closed limit sets of $\text{Prim}(A)$ was performed in [1]. We intend here to pursue the study begun in [2] of the topologies on the quotient of $\text{Prim}(A)$ determined by the algebra of the bounded scalar functions. We adopted a purely topological setting so no knowledge of the theory of C^* -algebras is needed for reading this paper. The method we chose is to substitute for the quotient of a possibly non-Hausdorff space X , via a homeomorphism, a quotient of a certain Hausdorff hyperspace of X .

In the following X denotes a locally compact space that is, X is a space in which every point has a neighbourhood base of compact sets. The algebra of all complex

1991 *Mathematics Subject Classification.* Primary: 54B15; Secondary: 54B20, 54D45, 46L05.
Key words and phrases. locally compact space, quotient space, Fell's topology.

valued bounded continuous functions on X , denoted $C^b(X)$, induces an equivalence relation on X : $x_1 \sim x_2$ if $f(x_1) = f(x_2)$ for every $f \in C^b(X)$. We let $\gamma(X)$ be the quotient space of this equivalence relation; the quotient map $q : X \longrightarrow \gamma(X)$ was called in [3, section III.3] the complete regularization of X and this construction was discussed in [2] for the special case of the primitive ideal space of a C^* -algebra. As in [2], we endow $\gamma(X)$ with the topology τ_{cr} , the weak topology defined by the bounded continuous functions on X viewed as functions on the quotient. Another natural topology on $\gamma(X)$ is the quotient topology τ_q . Always $\tau_{cr} \subset \tau_q$; it was shown in [2] that if X is compact or if q is open for τ_q or τ_{cr} then $\tau_q = \tau_{cr}$. The question whether these topologies are equal for any locally compact space was left open there. Of course, $C^b(\gamma(X))$ is always the same for both topologies. We shall prove in Section 2 that if X is σ -compact then $\tau_{cr} = \tau_q$. Recently D. W. B. Somerset found an example of a locally compact space X for which these two topologies are different. The example appears in an appendix to this paper and only its last paragraph, where it is shown that the topological space constructed there is homeomorphic to the primitive ideal space of a C^* -algebra, needs a minimum knowledge of operator algebra theory.

In Section 3 we discuss the class of second countable locally compact spaces X for which $\gamma(X)$ is a second countable locally compact Hausdorff space. Such spaces when serving as primitive ideal spaces of C^* -algebras were considered in [7] and [8]. We give a characterization of these spaces by using the tools developed in Section 2.

The family of all closed subsets of X will be denoted by $\mathcal{F}(X)$ and its subfamily that consists of all the nonempty closed subsets of X will be denoted $\mathcal{F}'(X)$. We shall equip $\mathcal{F}(X)$ with two topologies: the Fell topology, denoted here τ_s , that was defined in [9], and the lower semifinite topology of Michael, which we denote τ_w , see [12]. A base for τ_s consists of the family of all the sets

$$\mathcal{U}(C, \Phi) := \{S \in \mathcal{F}(X) \mid S \cap C = \emptyset, S \cap O \neq \emptyset, O \in \Phi\}$$

where C is a compact subset of X and Φ is a finite family of open subsets of X . The hyperspace $(\mathcal{F}(X), \tau_s)$ is always compact Hausdorff ([9, Lemma 1 and Theorem 1]). If X is second countable then this hyperspace is metrizable ([4, Lemme 2]). The family of all the sets $\mathcal{U}(\emptyset, \Phi)$ is a base for τ_w . The map η_X given by $\eta_X(x) := \overline{\{x\}}$

from X to $\mathcal{F}(X)$ is τ_w -continuous; it is not τ_s -continuous in general. It is one-to-one exactly when X is a T_0 space.

A subset S of X is called a limit subset if there is a net in X that converges to all the points of S . The family of all the closed limit subsets of X will be denoted $\mathcal{L}(X)$; it is a compact Hausdorff space with its relative τ_s -topology, metrizable if X is second countable, see [4, Théorème 12 and Lemme 2]. We put $\mathcal{L}'(X) := \mathcal{L}(X) \setminus \{\emptyset\}$. Then $(\mathcal{L}'(X), \tau_s)$ is a locally compact Hausdorff space; if X is compact then one easily sees that \emptyset is an isolated point of $(\mathcal{L}(X), \tau_s)$ hence $(\mathcal{L}'(X), \tau_s)$ is compact. The family of all (closed) maximal limit subsets is denoted $\mathcal{ML}(X)$ and $\mathcal{ML}^s(X)$ stands for its τ_s -closure in $\mathcal{L}'(X)$.

2. EQUIVALENCE RELATIONS ON $\mathcal{L}'(X)$

We define on $\mathcal{L}'(X)$ an equivalence relation: we say that $A \underset{1}{\sim} B$ if there is a finite sequence $\{F_i \mid 0 \leq i \leq n\}$ in $\mathcal{L}'(X)$ with $F_0 = A$ and $F_n = B$ such that $F_i \cap F_{i+1} \neq \emptyset$, $0 \leq i \leq n-1$. Let now $C_{\sim}^b(\mathcal{L}'(X))$ be the algebra of all \mathbb{C} -valued bounded τ_s -continuous functions on $\mathcal{L}'(X)$ that are constant on the equivalence classes with respect to $\underset{1}{\sim}$. For $A, B \in \mathcal{L}'(X)$ we shall say that $A \underset{2}{\sim} B$ if $f(A) = f(B)$ for every $f \in C_{\sim}^b(\mathcal{L}'(X))$. Then $\underset{2}{\sim}$ is an equivalence relation on $\mathcal{L}'(X)$ and by definition $A \underset{1}{\sim} B$ implies $A \underset{2}{\sim} B$. On $Q(X) := \mathcal{L}'(X) / \underset{2}{\sim}$ we shall consider two topologies: the quotient topology, τ_Q , defined by the quotient map $Q : \mathcal{L}'(X) \longrightarrow Q(X)$ when $\mathcal{L}'(X)$ is endowed with the τ_s topology and the completely regular topology τ_{CR} given by the functions of $C_{\sim}^b(\mathcal{L}'(X))$ considered as functions on $Q(X)$. Obviously $\tau_{CR} \subset \tau_Q$; it will follow from subsequent results that the question of equality between these two topologies parallels the situation between $(\gamma(X), \tau_{cr})$ and $(\gamma(X), \tau_q)$.

Let $f \in C^b(X)$; then f is constant on every closed limit subset of X . Define $f^{\mathcal{L}}$ on $\mathcal{L}'(X)$ by $f^{\mathcal{L}}(S) = f(x)$ where x is any point of $S \in \mathcal{L}'(X)$. Then $f^{\mathcal{L}}$ is τ_w -continuous of $\mathcal{L}'(X)$, thus $f^{\mathcal{L}} \in C_{\sim}^b(\mathcal{L}'(X))$. Indeed, if D is an open subset of \mathbb{C} then $U := \{x \in X \mid f(x) \in D\}$ is open hence $\{S \in \mathcal{L}'(X) \mid f^{\mathcal{L}}(S) \in D\} = \{S \in \mathcal{L}'(X) \mid S \cap U \neq \emptyset\}$ is in τ_w .

We have a converse to the statement about the continuity of $f^{\mathcal{L}}$ but first we need a lemma about τ_s -convergence in $\mathcal{L}'(X)$. It is included in [16, Lemma H.2] but we give below its simple proof for the sake of self sufficiency.

Lemma 2.1. *Let $\{S_\alpha \mid \alpha \in \mathcal{A}\}$ be a net in $\mathcal{L}'(X)$ that τ_s -converges to $S \in \mathcal{L}(X)$. If $x_\alpha \in S_\alpha$ for $\alpha \in \mathcal{A}$ and $\{x_\alpha\}$ converges to $x \in X$ then $x \in S$.*

Proof. Assuming that $x \notin S$ we let K be a compact neighbourhood of x disjoint from S . Thus $S \in \{T \in \mathcal{L}(X) \mid T \cap K = \emptyset\}$, hence eventually $S_\alpha \cap K = \emptyset$ and $\{x_\alpha\}$ cannot converge to x , a contradiction. \square

Theorem 2.2. *The map $f \longrightarrow f^\mathcal{L}$ is an isomorphism of $C^b(X)$ onto $C_\sim^b(\mathcal{L}'(X))$.*

Proof. We have to prove only the surjectivity of the map. Let $g \in C_\sim^b(\mathcal{L}'(X))$ and define f on X by $f(x) = g(S)$, S being any element in $\mathcal{L}'(X)$ such that $x \in S$, for instance the closure of $\{x\}$. Then f is well defined and we are going to show that it is continuous. Once this will be done we clearly shall have $f^\mathcal{L} = g$ and the proof will be finished. Let D be an open subset of \mathbb{C} and $x \in f^{-1}(D)$. We claim that there is a neighbourhood of x contained in $f^{-1}(D)$. If not then there is a net $\{x_\alpha \mid \alpha \in \mathcal{A}\}$ that converges to x but $f(x_\alpha) \notin D$ for every $\alpha \in \mathcal{A}$. We choose $S_\alpha \in \mathcal{L}'(X)$ such that $x_\alpha \in S_\alpha$ so that $f(x_\alpha) = g(S_\alpha)$ for each $\alpha \in \mathcal{A}$. The net $\{S_\alpha\}$ has a subnet $\{S_{\alpha'}\}$ that τ_s -converges to some $S \in \mathcal{L}(X)$. By Lemma 2.1 $x \in S$ hence $S \neq \emptyset$ and $g(S) = f(x) \in D$. The continuity of g implies that eventually $f(x_{\alpha'}) = g(S_{\alpha'}) \in D$ contradicting our choice of the net $\{x_\alpha\}$. \square

Remark 2.3. It follows from the τ_w -continuity of η_X , the definition of $f^\mathcal{L}$ for $f \in C^b(X)$ and the preceding proof that in the definition of $C_\sim^b(\mathcal{L}'(X))$ we can substitute τ_w -continuity for τ_s -continuity.

We can now define a one-to-one map χ from $\gamma(X)$ onto $Q(X)$ as follows: for $x \in X$ we let $\chi(q(x)) := Q(S)$ where S is any closed limit set that contains x . It readily follows from Theorem 2.2 that the map is well defined and it has the stated properties. As a direct consequence of the definitions we have for every $g \in C^b(\gamma(X))$ that $(g \circ q)^\mathcal{L} = g \circ \chi^{-1} \circ Q$. It is clear that χ is a homeomorphism of $(\gamma(X), \tau_{cr})$ onto $(Q(X), \tau_{CR})$.

Proposition 2.4. *The map χ defined above is a homeomorphism from $(\gamma(X), \tau_q)$ onto $(Q(X), \tau_Q)$.*

Proof. Let $O \subset Q(X)$ be open in the quotient topology. Then $U := q^{-1}(\chi^{-1}(O))$ is the union of all the elements of $Q^{-1}(O)$ and we claim that U is open. Otherwise, there are $x \in U$ and a net $\{x_\alpha\}$ in $X \setminus U$ that converges to x . For each index α we choose S_α such that $x_\alpha \in S_\alpha$. This compels each S_α to belong to the closed subset $\mathcal{L}'(X) \setminus Q^{-1}(O)$ of $\mathcal{L}'(X)$. By passing to a subnet if necessary we may suppose that $\{S_\alpha\}$ τ_s -converges to some $S \in \mathcal{L}(X)$. Lemma 2.1 yields $x \in S$ thus $S \in Q^{-1}(O)$. On the other hand, $S \notin Q^{-1}(O)$ as the τ_s -limit of S_α , a contradiction. Therefore U is an open subset of X , $\chi^{-1}(O)$ is a τ_q -open subset of $\gamma(X)$ and the continuity of χ is established.

Suppose now that $V \subset \gamma(X)$ is τ_q -open. Then

$$Q^{-1}(\chi(V)) = \{S \in \mathcal{L}'(X) \mid S \subset q^{-1}(V)\} = \{S \in \mathcal{L}'(X) \mid S \cap q^{-1}(V) \neq \emptyset\}$$

is τ_w -open hence τ_s -open. Thus $\chi(V)$ is τ_Q -open. \square

Before stating the main result of this section we need a lemma.

Lemma 2.5. *If X is σ -compact then $(\mathcal{L}'(X), \tau_s)$ is σ -compact too.*

Proof. Let $X = \cup_{n=1}^{\infty} K_n$ where each K_n is compact and set $L_n = \{S \in \mathcal{L}'(X) \mid S \cap K_n \neq \emptyset\}$. Clearly $\mathcal{L}'(X) = \cup_{n=1}^{\infty} L_n$ and we are going to show that each L_n is τ_s -compact. Let $\{S_\alpha\}$ be a net in L_n and without loss of generality we shall suppose that it τ_s -converges to some $S \in \mathcal{L}(X)$. For each α choose $x_\alpha \in S_\alpha \cap K_n$. By passing to a subnet we may suppose that $\{x_\alpha\}$ converges to some $x \in K_n$. Lemma 2.1 yields $x \in S$ thus $S \in L_n$. \square

Theorem 2.6. *If X is σ -compact then $\tau_{CR} = \tau_q$ and $\gamma(X)$ is paracompact.*

Proof. In view of Proposition 2.4 and the remarks preceding its statement it will suffice to show that $\tau_{CR} = \tau_Q$ and that $Q(X)$ is paracompact. Now, $\mathcal{L}'(X)$ with the τ_s topology is locally compact Hausdorff and σ -compact by Lemma 2.5 hence Lindelöf. Its quotient space $(Q(X), \tau_Q)$ is Hausdorff since the real valued bounded continuous functions on $Q(X)$ separate its points. It follows from Theorem 1 of [14] that $Q(X)$ with its τ_Q topology is a paracompact space. In particular it is also completely regular, hence $\tau_{CR} = \tau_Q$ and we are done.

□

Remark 2.7. In all of the above we could have used the space $\mathcal{ML}^s(X)$ instead of $\mathcal{L}'(X)$.

We are going to treat another situation when the two topologies on $\gamma(X)$ coincide but first we have to introduce a new relation on the space X that was considered in [2] for the primitive ideal space of a C^* -algebra. For $x, y \in X$ we shall write $x \underset{H}{\sim} y$ if x and y cannot be separated by disjoint open subsets of X . This is the same as saying that there is a closed limit subset of X to which both x and y belong. In general this is not a transitive relation. Clearly, if $\underset{H}{\sim}$ is an equivalence relation then each equivalence class for it is the union of all the elements in an equivalence class with respect to $\underset{1}{\sim}$ on $\mathcal{L}'(X)$ and each such union is an equivalence class for $\underset{H}{\sim}$. The following result is the same as Proposition 3.2 of [2] when X is the primitive ideal space of a C^* -algebra but the proof below differs in part from that given there.

Proposition 2.8. *Suppose $\underset{H}{\sim}$ is an open equivalence relation. Then each equivalence class is a maximal limit set and each maximal limit set of X is an equivalence class for $\underset{H}{\sim}$. The relations $\underset{H}{\sim}$ and $\underset{H}{\sim}$ are the same, $\tau_q = \tau_{cr}$, the quotient map q is open and $\gamma(X)$ is a locally compact Hausdorff space.*

Proof. Let S be an equivalence class for $\underset{H}{\sim}$. We are going to show that for every finite set $\{x_i \mid 1 \leq i \leq n\} \subset S$ and every neighbourhood V_i of x_i , $1 \leq i \leq n$, we have $\cap_{i=1}^n V_i \neq \emptyset$; by [4, Lemme 9] this will imply that S is a limit set. The claim is obviously valid for any pair of points of S . Suppose that it is true for any subset of $n-1$ points of S and let $x_i \in S$ with an arbitrary neighbourhood V_i , $1 \leq i \leq n$. Let U be the open saturation of V_n for $\underset{H}{\sim}$. Then $x_i \in U$ and $U_i := V_i \cap U$ is a neighbourhood of x_i , $1 \leq i \leq n-1$. By the induction hypothesis $W := \cap_{i=1}^{n-1} U_i$ is a non-void open set. Choose $x \in W$; there is $y \in V_n$ such that $x \underset{H}{\sim} y$. By the definition of $\underset{H}{\sim}$ we have $\cap_{i=1}^n V_i \supset W \cap V_n \neq \emptyset$ and the claim is established. Now if S' is a limit set with $S' \supset S$ then each point of S' is $\underset{H}{\sim}$ -equivalent to each point of S hence $S' = S$. Thus S is a maximal limit set. Since all the points of a maximal limit set are $\underset{H}{\sim}$ -equivalent no two different maximal limit sets can intersect and each maximal limit set is an equivalence class for $\underset{H}{\sim}$.

From here on we follow the proof of [2, Proposition 3.2]. If S_1 and S_2 are two different \sim_H -classes and $x_i \in S_i$ then x_1 and x_2 have two disjoint open neighbourhoods V_1 and V_2 , respectively. No point of V_1 can be \sim_H -equivalent to any point of V_2 because $V_1 \cap V_2 = \emptyset$. Hence the open quotient map of X onto X/\sim_H maps V_1 and V_2 onto two disjoint neighbourhoods of S_1 and S_2 respectively, which means that the quotient space is Hausdorff. Since the quotient map is open the quotient space is also locally compact. Clearly if $x \sim_H y$ then $x \sim y$. By the complete regularity of X/\sim_H we conclude that if x and y are not \sim_H -equivalent they are also not \sim -equivalent. Thus \sim_H and \sim are identical. Since $(\gamma(X), \tau_q)$ and $(\gamma(X), \tau_{cr})$ have the same bounded continuous functions and both are completely regular, the identity map is a homeomorphism. \square

3. \mathcal{CR} -SPACES

In this section we shall discuss a class of locally compact second countable spaces. A locally compact second countable space has a countable base consisting of interiors of compact subsets since the family of the interiors of all the compact subsets is a base and as such it must contain a countable base by [10, Problem 1.F]. Thus such a space X is σ -compact and by Theorem 2.6 we have only one topology on $\gamma(X)$ that will be of interest for us. Of course, this is true also for $Q(X)$.

As we remarked in the previous section, the real valued bounded continuous functions on $\gamma(X)$ separate the points of this space. It turns out that when X is second countable a countable family of such functions will suffice.

Proposition 3.1. *If X is second countable then there is a countable family of real valued bounded continuous functions on $\gamma(X)$ that separates the points of $\gamma(X)$.*

Proof. From Proposition 2.4 we gather that it will be enough to show that such a countable family of functions exists on $Q(X)$. Recall that $(\mathcal{L}'(X), \tau_s)$ is locally compact Hausdorff and second countable, in particular σ -compact. Let $\{K_n\}$ be an increasing sequence of τ_s -compact subsets that covers $\mathcal{L}'(X)$. Then $C(K_n)$, the algebra of all real continuous functions on the compact metrizable space K_n is separable. Hence $C_\sim^b(\mathcal{L}'(X))|_{K_n} \subset C(K_n)$ is also separable. Thus there is a countable family $\{f_n^m \mid 1 \leq m < \infty\} \subset C_\sim^b(\mathcal{L}'(X))$ such that $\{f_n^m|_{K_n} \mid 1 \leq m < \infty\}$ is dense

in $C_\sim^b(\mathcal{L}'(X)) \mid_{K_n}$. Now, if g and h are real bounded continuous functions on $Q(X)$ then

$$\sup \{|g(y) - h(y)| \mid y \in Q(K_n)\} = \sup \{|g \circ Q(S) - h \circ Q(S)| \mid S \in K_n\}.$$

Hence, viewing the elements of $C_\sim^b(\mathcal{L}'(X))$ as functions on $Q(X)$, the family $\{f_n^m \mid 1 \leq m < \infty\}$ separates the points of $Q(K_n)$ and $\{f_n^m \mid 1 \leq m, n < \infty\}$ separates the points of $Q(X)$.

□

Definition 3.2. A second countable locally compact space X will be called a \mathcal{CR} -space if $\gamma(X)$ is a second countable locally compact Hausdorff space.

In [7] a separable C^* -algebra whose primitive ideal space is a \mathcal{CR} -space in the above terminology was called a \mathcal{CR} -algebra. There the conditions imposed on the quotient of the primitive ideal space were for the topology τ_{cr} . Not every separable C^* -algebra is a \mathcal{CR} -algebra thus not every second countable locally compact space is a \mathcal{CR} -space. An example is given in [3, Example 9.2]. The class of \mathcal{CR} -algebras was found in [7] and [8] useful for the study of certain C^* -dynamical systems and the corresponding crossed products. It was remarked in [8] that for a second countable locally compact space X each of the following properties is sufficient to ensure that it is a \mathcal{CR} -space: X is Hausdorff, X is compact, \sim_H is an open equivalence relation on X . Of course, this was done in [8] only for the primitive ideal space of a C^* -algebra so we shall reproduce and adapt the arguments for the general situation. The case of a Hausdorff space is trivial. If X is compact then $\gamma(X)$ is compact too. Recall that $\mathcal{L}'(X)$ is τ_s -compact. Theorem 2.2 and the definition of the quotient topology on $\gamma(X)$ yield an isomorphism of the algebra $C(\gamma(X))$ into the separable algebra $C(\mathcal{L}'(X))$ hence $C(\gamma(X))$ and $\gamma(X)$ is second countable. Alternatively, we can use the Proposition 3.1 to infer that $\gamma(X)$ is metrizable. Now suppose that X is a second countable locally compact space for which \sim_H is an open equivalence relation. Then by Proposition 2.8, $\gamma(X)$ is locally compact Hausdorff. Since the quotient map is continuous and open it is easily seen that $\gamma(X)$ is second countable. We give below a characterization of the \mathcal{CR} -spaces.

Theorem 3.3. *Let X be a second countable locally compact space. The following conditions are equivalent:*

- (i) X is a \mathcal{CR} -space;
- (ii) $\gamma(X)$ is locally compact;
- (iii) $\gamma(X)$ is first countable.

Proof. (i) \Rightarrow (ii). This is immediate.

(ii) \Rightarrow (iii). By assumption $\gamma(X)$ is locally compact Hausdorff and σ -compact since X is σ -compact. Then, by [6, Theorem 7.2], there is an increasing sequence of open sets $\{U_n\}$ in $\gamma(X)$ such that $\gamma(X) = \cup_{n=1}^{\infty} U_n$, \overline{U}_n is compact and $\overline{U}_n \subset U_{n+1}$ for every n . Proposition 3.1 yields a sequence $\{g_k\}$ of continuous functions from $\gamma(X)$ to the interval $[0, 1]$ that separates the points of $\gamma(X)$. The restrictions of the functions $\{g_k\}$ to the compact Hausdorff space \overline{U}_n allow us to define a homeomorphism of \overline{U}_n into $[0, 1]^{\aleph_0}$. Hence \overline{U}_n is metrizable and each open set U_n is second countable. We conclude that $\gamma(X)$ is second countable. We actually proved that (i) follows from (ii) which is apparently more than we needed.

(iii) \Rightarrow (i). From Proposition 2.4 and the hypothesis it follows that $Q(X)$ is first countable. We know that it is a Hausdorff space. It has been noted above that $\mathcal{L}'(X)$ is locally compact Hausdorff and under the present hypothesis on X it is also second countable. Thus $Q(X)$ is a Hausdorff quotient of a second countable locally compact Hausdorff space. Then [15, Theorem 3] implies that $Q(X)$ is locally compact and second countable and the same properties are shared by $\gamma(X)$ by Proposition 2.4.

□

Another characterization of \mathcal{CR} -spaces can be given in terms of the quotient map q . A continuous map φ from a topological space Y onto a topological space Z was called in [13] a bi-quotient map if for every $z \in Z$ and every open cover of $\varphi^{-1}(z)$ there are finitely many sets $\{U_i\}$ in the cover such that the interior of $\cup_{i=1}^n \varphi(U_i)$ is a neighbourhood of z . It follows from [13, Proposition 3.3(d) and Proposition 3.4] that whenever φ is a quotient map of the second countable locally compact space Y onto the Hausdorff space Z then Z is locally compact and second countable if and only if φ is bi-quotient. If we adapt this general result to our situation we get

Proposition 3.4. *The second countable locally compact space X is a CR -space if and only if the quotient map q is bi-quotient.*

REFERENCES

1. R. J. Archbold, *Topologies for primal ideals*, J. London Math. Soc. **36** (1987), 524–542.
2. R. J. Archbold and D. W. B. Somerset, *Quasi-standard C^* -algebras*, Math. Proc. Cambridge Philos. Soc. **107** (1990), 349–360.
3. J. D. Dauns and K. H. Hofmann, *Representation of rings by sections*, Mem. Amer. Math. Soc. **83** (1968), 1–180.
4. J. Dixmier, *Sur les espaces localement quasi-compact*, Canad. J. Math. **20** (1968), 1093–1100.
5. J. Dixmier, *C^* -algebras*, North-Holland, Amsterdam, 1977.
6. J. Dugundji, *Topology*, Allyn and Bacon, Boston, 1966.
7. S. Echterhoff and D. P. Williams, *Crossed products by $C_0(X)$ -actions*, J. Funct. Anal. **158** (1998), 113–151.
8. S. Echterhoff and D. P. Williams, *Locally inner actions on $C_0(X)$ -algebras*, J. Operator Theory **45** (2001), 131–160.
9. J. M. G. Fell, *A Hausdorff topology for the closed sets of a locally compact non-Hausdorff space*, Proc. Amer. Math. Soc. **13** (1962), 472–476.
10. J. L. Kelley, *General Topology*, Van Nostrand, Princeton, NJ, 1955.
11. A. J. Lazar, *Hyperspaces of closed limit sets*, to appear.
12. E. Michael, *Topologies on spaces of subsets*, Trans. Amer. Math. Soc. **71** (1951), 152–182.
13. E. Michael, *Bi-quotient maps and cartesian products of quotient maps*, Ann. Inst. Fourier **18** (1968), 287–302.
14. K. Morita, *On decomposition spaces of locally compact spaces*, Proc. Japan Acad. **32** (1956), 544–548.
15. A. H. Stone, *Metrisability of decomposition spaces*, Proc. Amer. Math. Soc. **7** (1956), 690–700.
16. D. P. Williams, *Crossed products of C^* -algebras*, Mathematical Surveys and Monographs, vol. 134, American Mathematical Society, Providence, RI, 2007.

SCHOOL OF MATHEMATICAL SCIENCES, TEL AVIV UNIVERSITY, TEL AVIV 69778, ISRAEL
E-mail address: aldo@post.tau.ac.il